



This information is current as of August 19, 2013.

MHC Class I and TCR Avidity Control the CD8 T Cell Response to IL-15/IL-15R α Complex

Thomas A. Stoklasek, Sara L. Colpitts, Henry M. Smilowitz and Leo Lefrançois

J Immunol 2010; 185:6857-6865; Prepublished online 1 November 2010; doi: 10.4049/jimmunol.1001601 http://www.jimmunol.org/content/185/11/6857

Supplementary http://www.jimmunol.org/content/suppl/2010/11/01/jimmunol.100160 Material 1.DC1.html References This article cites 48 articles, 36 of which you can access for free at: http://www.jimmunol.org/content/185/11/6857.full#ref-list-1 **Subscriptions** Information about subscribing to *The Journal of Immunology* is online at: http://jimmunol.org/subscriptions Permissions Submit copyright permission requests at: http://www.aai.org/ji/copyright.html **Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at: http://jimmunol.org/cgi/alerts/etoc





MHC Class I and TCR Avidity Control the CD8 T Cell Response to IL-15/IL-15Rα Complex

Thomas A. Stoklasek,* Sara L. Colpitts,* Henry M. Smilowitz,^{†,‡} and Leo Lefrançois*

IL-15 operates via a unique mechanism termed transpresentation. In this system, IL-15 produced by one cell type is bound to IL-15R α expressed by the same cell and is presented to apposing cells expressing the IL-15R β/γ C complex. We have shown that administering soluble IL-15R α complexed with IL-15 can greatly enhance IL-15 activity. We now show that the naive CD8 T cell response to exogenous IL-15/IL-15R α complex is MHC class I dependent. In the absence of β 2 microglobulin, naive CD8 T cells scarcely proliferated in response to IL-15/IL-15R α complex, whereas memory cells proliferated, although to a lesser extent, compared with levels in control mice. The loss of β 2m or FcRn slightly reduced the extended half-life of IL-15/IL-15R α complex. In addition, we demonstrated a link between TCR avidity and the ability of a T cell to respond to IL-15/IL-15R α complex. Thus, T cells expressing low-avidity TCR responded poorly to IL-15/IL-15R α complex, which correlated with a poor homeostatic proliferative response to lymphopenia. The inclusion of cognate peptide along with complex resulted in enhanced proliferation, even when TCR avidity was low. IL-15/IL-15R α complex treatment, along with peptide immunization, also enhanced activation and the migratory ability of responding T cells. These data suggest that IL-15/IL-15R α complex has selective effects on Ag-activated CD8 T cells. Our findings have important implications for directing IL-15/IL-15R α complex has selective effects.

verlapping factors regulate the naive and memory CD8 T cell pools in the periphery. Naive CD8 T cells, but not memory CD8 T cells, require contact with self-peptide presented by MHC class I molecules, whereas both populations require IL-7 for their survival (1-5). Furthermore, memory CD8 T cells require IL-15 for their continued homeostatic proliferation in an immunosufficient background (6). $IL15^{-/-}$ and $IL-15R\alpha^{-/-}$ mice have a significantly smaller naive CD8 T cell compartment, possibly due to reduced CD8 T cell thymic development and survival in the periphery (7). These requirements for maintenance and survival are paralleled when T cells are placed in an immunodeficient environment. Specifically, naive CD8 T cells require host MHC class I and IL-7 expression to undergo acute homeostatic proliferation in irradiated or RAG-deficient hosts (5, 8-11). IL-15 is not required for initiating naive CD8 T cell acute homeostatic proliferation, but rather it helps to sustain the process at later time points (12). In contrast, memory CD8 T cells proliferate in the absence of MHC class I, but they require IL-7 and IL-15 for their maximum proliferation in a lymphopenic setting (3, 5).

The identification of the mechanism of IL-15–mediated activity, namely transpresentation (13, 14), led to the use of precomplexed IL-15 and soluble IL-15R α as immunostimulants (15–17). Thus, presentation of rIL-15 bound with high affinity to soluble rIL-

Received for publication May 14, 2010. Accepted for publication October 1, 2010.

 $15R\alpha$ drives robust activation of cells expressing IL-2/15R β and the γ C in vitro and in vivo. Memory CD8 T cells and NK cells respond most vigorously, and IL-15/IL-15R α (IL-15 complex) treatment augments tumor clearance (16, 18, 19).

Surprisingly, naive CD8 T cells also respond to IL-15 complex, despite their low expression level of IL-15R β (11, 16). Of note, the naive CD8 T cell response to IL-15 complex mimics the response to cognate Ag in that the naive CD8 T cell population undergoes an expansion phase and acquires an activated phenotype and effector functions, including the ability to secrete IFN- γ and mediate Ag-specific cytolytic function. Importantly, no exogenous Ag was required to initiate this response. Similarly, naive CD8 T cells undergoing acute homeostatic proliferation in immunodeficient hosts acquire a memory phenotype with increased expression of CD44, IL-2/IL-15RB and YC, Ly6C, and CD69 (10, 20, 21). Some initial studies stated that naive CD8 T cells proliferating in a lymphopenic environment do not acquire effector function after transfer (22), whereas others showed the acquisition of potent effector responses (20, 21). However, this phenomenon requires multiple rounds of division and develops relatively late posttransfer, which may explain the differences between these studies. In addition, acute homeostatic proliferation results in the appearance of long-term stable memory-phenotype CD8 T cells (23, 24). Thus, acute lymphopenic proliferation and IL-15/IL- $15R\alpha$ -induced proliferation of naive CD8 T cells result in the acquisition of memory phenotypes, although IL-15 complex stimulation operates with more rapid induction of proliferation (16).

Although acute slow homeostatic proliferation occurs without the addition of cognate Ag, the process is dependent on the presence of MHC class I presenting self-peptide (8, 9, 25). However, exogenous peptides derived from normal flora drive a subset of naive polyclonal T cells to undergo rapid homeostatic proliferation (26). Furthermore, although some naive TCR transgenic T cells, such as SIINFEKL-specific OT-I TCR transgenic CD8 T cells, undergo massive proliferation when placed in a lymphopenic environment, this is not the case with all naive CD8 T cells (9–11). This effect

^{*}Department of Immunology, Center for Integrated Immunology and Vaccine Research, [†]Department of Diagnostic Imaging and Therapeutics, and [‡]Department of Cell Biology, University of Connecticut Health Center, Farmington, CT 06030

Address correspondence and reprint requests to Dr. Leo Lefrançois, Department of Immunology, University of Connecticut Health Center, 263 Farmington Avenue, Farmington, CT 06107. E-mail address: llefranc@neuron.uchc.edu

The online version of this article contains supplemental material.

Abbreviations used in this paper: hIL-15, human IL-15; IL-2 complex, IL- 2α /IL-2; peptide/IL-15 complex, SIINFEKL peptide and IL-15 complex; pLN, peripheral lymph node; VSV-OVA, vesicular stomatitis virus expressing OVA.

Copyright © 2010 by The American Association of Immunologists, Inc. 0022-1767/10/\$16.00

seems to be due, in part, to differences in TCR triggering, whether due to TCR density, negative-regulator CD5 levels, TCR avidity, or TCR promiscuity (8, 9, 27–29). These findings indicate that the strength of the TCR stimulus is directly related to the extent of proliferation in response to lymphopenia (10).

Because naive CD8 T cells responding to IL-15 complex seem to undergo a similar activation process as cells responding to lymphopenia (16, 21), we set out to compare the requirements for induction of T cell activation by these conditions. The results indicated that IL-15 complex-driven CD8 T cell activation operates through TCR signaling and not solely through cytokine receptors. Understanding the mechanism through which IL-15 complex modulates the immune system holds implications for the proper application of IL-15 immunotherapy.

Materials and Methods

Mice

C57BL/6-CD45.2 mice were purchased from The Jackson Laboratory (Bar Harbor, ME) and the Charles River-National Cancer Institute. C57BL/6-CD45.1 mice were purchased from Charles River-National Cancer Institute (Frederick, MD). B10.D2 mice (recipients of C4 transfers) were purchased from The Jackson Laboratory. The OT-I mouse line was maintained as a C57BL/6-CD45.1 line on a $Rag^{-/-}$ background. FcRn^{-/-} mice were a generous gift from Dr. Lynn Puddington (University of Connecticut Health Center) (30). β 2m^{-/-} mice and their respective C57BL/6-CD45.2 controls were purchased from Taconic Farms (Germantown, NY). The F5 (31) and C4 (32) mice were generously provided by Dr. Linda Cauley and Dr. Adam Adler, respectively (University of Connecticut Health Center). All procedures were carried out under National Institutes of Health Genter Animal Care Committee.

IL-15 treatment

Human IL-15 (hIL-15) was generously provided by Amgen (Thousand Oaks, CA). Mouse rIL-15R α -Fc chimeric molecule was purchased from R&D Systems (Minneapolis, MN). hIL-15 and mouse rIL-15R α -Fc, both suspended in PBS, were mixed and incubated for 30 min at 37°C. Each mouse, unless specifically noted, received 2.5 µg IL-15 alone or precomplexed with 15 µg mouse rIL-15R α -Fc in 200 µl PBS i.p.

Purification and adoptive transfer of cells

Single-cell suspensions were created in HBSS by homogenizing spleens and lymph nodes using frosted glass slides. RBCs were lysed using Trisammonium chloride buffer, and cells were filtered through Nitex. CD44¹⁰ CD8 T cells (includes TCR transgenic T cells) were enriched using a combination of anti-CD44-biotin and antibiotin microbeads (Miltenyi Biotec, Auburn, CA), as previously described (33). In experiments comparing the proliferation of CD44^{lo} and CD44^{hi} polyclonal CD8 T cells, both populations were purified by flow cytometry using a FACSVantage SE (BD Biosciences, San Jose, CA). For CFSE labeling, cells were incubated for 10 min at 37°C with CFSE (0.01 mM; Invitrogen, Carlsbad, CA), the reaction was quenched with HBSS with 5% FCS (34), and the cells were washed twice. CFSE-labeled cells were resuspended in PBS and injected i.v. into congenic mice. Unless otherwise specified in the figure legends, the number of cells transferred was between 1×10^5 and 3×10^5 . For infections, 24 h post-OT-I transfer mice were infected i.v. with 1×10^5 vesicular stomatitis virus expressing OVA (VSV-OVA).

Isolation of cells from tissues

Using frosted glass slides, spleen and lymph nodes were homogenized into single-cell suspensions in HBSS/5% bovine serum. Lungs, liver, and brains were minced into small pieces and incubated with collagenase buffer (RPMI 1640, 2% FCS, containing HEPES, pen-strep, glutamine, gentamicin, CaCl₂, MgCl₂, and collagenase [100 U/ml; Invitrogen, Carlsbad, CA]) for 1 h (lung and liver) or 30 min (brain) at 37°C under agitation. At the end of incubation, remaining tissue pieces were crushed through 70- μ m nylon mesh filter (BD Biosciences). The resulting cells were pelleted and resuspended in 44% Percoll buffer (GE Life Sciences, Piscataway, NJ) and underlain with 67% Percoll buffer. Percoll gradients were extracted, washed, and resuspended in HBSS/5% bovine serum/HEPES, pen-strep, glutamine, gentamicin.

Flow-cytometric analysis

Cells were isolated at the indicated times and analyzed for the presence of donor cells using CD45 or Thy1 allele status and their expression of surface markers and CFSE intensity. For staining, lymphocytes were suspended in PBS/0.2% BSA/0.1% NaN₃ (FACS buffer) at a concentration of $5 \times 10^6/100 \,\mu$ l and stained with indicated Abs at 4°C for 20 min, washed, and fixed in PBS with 2% paraformaldehyde. Samples were analyzed using FlowJo Software (Tree Star, Ashland, OR).

Peptide activation and treatments

Congenic naive T cells were transferred into appropriate hosts (OT-I and F5 into B6 hosts) that were subsequently treated as described in the figure legends. For peptide treatments, mice received 100 μ g of the respective peptide, unless stated otherwise. The peptide sequences were as follows from N terminus to C terminus: SIINFEKL (OT-I-specific epitope presented by H-2K^b) and ASNENMETM (F5-specific epitope from influenza virus NP presented by H-2D^b). For combination peptide and IL-15 complex treatment, peptides were added to the mixture of IL-15 and L-15R α -Fc (as described above) prior to injection. All treatments were given i.p.

Serial adoptive transfers and T cell-migration assay

We performed serial adoptive transfers to isolate OT-I T cells subjected to different activating conditions and to test their migratory potential. Naive CD44¹⁰ CD45.1 OT-I cells were transferred into naive CD45.2 B6 hosts; 1 d later, hosts were injected with IL-15 complex, a mixture of 100 µg SIINFEKL and IL-15 complex, or VSV-OVA. Four days posttreatment, spleens were harvested and processed into single-cell suspensions. Splenocytes were enriched for activated CD45.1 donor OT-I cells using MACS technology. In short, single-cell suspensions were stained with CD45.1 biotin and enriched for CD45.1 OT-I using antibiotin microbeads and MACS LS separation columns, according to Miltenyi Biotec protocols. Approximately 1×10^6 enriched activated CD45.1 OT-I cells, mixed with CD45.2 naive B6 splenocytes to equalize the total number of cells transferred into each host, were transferred into secondary CD45.2 B6 hosts. For the naive OT-I control, naive CD44¹⁰ OT-I cells were isolated as described above, but hosts were left untreated. Two days following secondary OT-I transfer, cells from spleen, peripheral lymph nodes (pLNs) (inguinal, brachial, axillary, and cervical lymph nodes), lung, and liver were isolated and processed as described above. Presence of CD45.1 donor OT-I cells was analyzed by flow cytometry. The percentage of donor cells in each tissue was formulated as follows: the number of CD45.1 donor OT-I cells in a particular tissue was divided by the total number of CD45.1 donor OT-I cells in the spleen, pLNs, liver, and lung and then multiplied by 100.

Results

Naive CD8 T cell response to IL-15 complex is MHC class I dependent

As we previously illustrated, naive CD8 T cells acquire an activated phenotype and effector function in response to IL-15 complex (16). Although cognate Ag is not required for these effects, it is not known whether the response is MHC class I dependent. To test this possibility, we transferred CFSE-labeled CD44^{lo} CD45.1 naive OT-I TCR transgenic CD8 T cells (Fig. 1A) into sublethally irradiated (500 rad) CD45.2 B6 and MHC class I-deficient ($\beta 2m^{-/-}$) mice on day -1. Hosts were irradiated to prevent rejection of donor cells in $\beta 2m^{-/-}$ mice. On day 0, B6 and $\beta 2m^{-/-}$ mice were treated with PBS or IL-15 complex (2.5 µg IL-15 and 15 µg sIL- $15R\alpha$ -Fc; doses used for all experiments), and splenocytes were analyzed on day 4 posttreatment for OT-I proliferation. As expected, the naive CD44^{lo} OT-I cells underwent homeostatic proliferation in irradiated B6 hosts, and this proliferation was substantially augmented with IL-15 complex treatment (Fig. 1A). In contrast, in untreated irradiated $\beta 2m^{-/-}$ hosts, naive OT-I CD8 T cells did not proliferate. With IL-15 complex treatment, only a low level of proliferation was induced in $\beta 2m^{-1}$ mice, as evidenced by blunted CFSE dilution and reduced cell numbers (Fig. 1A). The minor response detected in $\beta 2m^{-/-}$ hosts may be due to the interaction of OT-I cells with other MHC class I^+ donor cells or with MHC class II present in the host (29, 35).



FIGURE 1. Naive CD8 T cell response to IL-15 complex requires MHC class I. A, A total of 5×10^5 CFSE-labeled naive CD45.1 CD4410 OT-I TCR transgenic CD8 T cells was injected i.v. into CD45.2 B6 or $\beta 2m^{-/-}$ hosts that were sublethally irradiated (500 rad) 12 h earlier. One day later, mice were injected i.p. with PBS or 2.5 μ g hIL-15 + 15 μ g mIL-15R α -Fc. Four days after treatment, donor cell proliferation and accumulation were examined in the spleen. A total of 5×10^5 CFSE-labeled CD45.1 CD44^{lo} polyclonal CD8 T cells (B) and 3×10^5 CFSE-labeled Thy1.1 CD44^{hi} polyclonal CD8 T cells (C) were coinjected i.v. into CD45.2 Thy1.2 B6 and $\beta 2m^{-/-}$ mice that were sublethally irradiated (500 rad) 12 h earlier. One day later, mice were injected i.p. with PBS or 2.5 μg hIL-15 + 15 μ g mouse IL-15R α -Fc. Four days later, donor cell proliferation and accumulation were examined in the spleen. Data are representative of at least two experiments (n = 3).



Thus, the naive CD8 T cell response to IL-15 complex was largely dependent on the presence of MHC I.

Contrary to naive CD8 T cells, the proliferation of CD44^{hi} CD8 T cells in a lymphopenic environment is not MHC class I dependent (2, 3). To test whether this was also the case with IL-15 complex, sorted CFSE-labeled CD44^{hi} Thy1.1 and CD44^{lo} CD45.1 polyclonal CD8 T cells were transferred together into sublethally irradiated (550 rad) CD45.2/Thy1.2 B6 and $\beta 2m^{-/-}$ hosts on day -1. On day 0, B6 and $\beta 2m^{-\prime-}$ mice were treated with PBS or IL-15 complex, and splenocytes were analyzed for the proliferation of donor cells on day 4. Consistent with the OT-I data, the proliferation of CD44^{lo} polyclonal CD8 T cells was greatly reduced in $\beta 2m^{-\prime-}$ hosts compared with B6 hosts (Fig. 1B). In contrast, treatment with IL-15 complex substantially enhanced proliferation of memory-phenotype cells in B6 hosts and, to a lesser extent, in $\beta 2m^{-/-}$ hosts (Fig. 1*C*). The results indicated that the response of naive CD8 T cells to IL-15 complex had a greater dependence on MHC class I than did memory-phenotype cells, suggesting that complex treatment mimicked events seen in a lymphopenic situation (3, 8, 10).

FcRn partially regulates IL-15/IL-15Rα complex pharmacology

Although $\beta 2m^{-/-}$ mice lack MHC class I expression, $\beta 2m$ is also required for expression of the neonatal Fc receptor FcRn (36). FcRn plays an important role in IgG catabolism through IgG up-

take and recycling, such that FcRn deficiency results in reduced IgG half-life. Because the IL-15Rα component of IL-15 complex contains an IgG Fc region, we investigated whether B2m expression affected complex half-life. To this end, B6 and $\beta 2m^{-/-}$ hosts were administered equal amounts of IL-15 complex, and serum IL-15 levels were measured at various times after treatment by ELISA (16). At early time points, more IL-15 was detected in the serum of $\beta 2m^{-/-}$ mice compared with control mice (Fig. 2A). However, IL-15 decayed more rapidly in $\beta 2m^{-/-}$ mice, and the total amount of available IL-15 was ~60% of that in B6 mice, as measured by calculating the area under each curve. The role of FcRn in this phenomenon was directly assessed using FcRn^{-/-} hosts. IL-15 serum half-life exhibited similar kinetics in FcRn^{-/-} hosts as in $\beta 2m^{-/-}$ hosts, although no early spike in IL-15 levels was noted as in $\beta 2m^{-/-}$ mice (Fig. 2B). As seen in $\beta 2m^{-/-}$ mice, we noted that the total amount of available IL-15 in $FcRn^{-/-}$ mice was ~60% of that in B6 mice. We also measured the proliferation of naive OT-I cells in sublethally irradiated B6 and FcRn^{-/-} hosts. Without complex treatment, OT-I cells divided equally well in sublethally irradiated $FcRn^{-/-}$ and B6 hosts (Fig. 2C). However, OT-I cell proliferation was partially reduced in IL-15 complextreated FcRn^{-/-} mice compared with B6 hosts, which may be explained by the reduction in IL-15 half-life and total availability (Fig. 2B). Nonetheless, proliferation of naive OT-I cells in FcRn^{-/-} mice was substantially greater than that obtained in $\beta 2m^{-/-}$ mice (Fig. 1A). Thus, although the lack of FcRn



FIGURE 2. Naive CD8 T cell response to IL-15 complex is partially dependent on FcRn. A, B6 and $\beta 2M^{-/-}$ mice were injected i.p. with 2.5 µg hIL-15 + 15 μg mouse IL-15Rα-Fc, and hIL-15 was measured in the blood at various time points after injection (0, 0.5, 1, 2, 4, 8, 24, 48, 72, and 96 h) by ELISA. B, As in A, but using B6 and $FcRn^{-/-}$ mice and evaluating at ~0.25, 0.5, 1, 2, 4, 8, 24, 48, 72, and 96 h. C, A total of 4.0 \times 10 5 CFSE-labeled CD45.1 CD44 $^{\rm lo}$ OT-I CD8 T cells were injected i.v. into CD45.2 B6 or FcRn^{-/} mice sublethally irradiated (500 rad) 12 h earlier. One day later, mice were treated with PBS or 2.5 µg hIL- $15 + 15 \mu g$ mouse IL- $15R\alpha$ -Fc. Four days later, donor cell proliferation and accumulation were examined in the spleen. Data are representative of at least two experiments (n = 2-4).

influenced the naive CD8 T cell response in $\beta 2m^{-/-}$ mice, the majority of IL-15 complex-driven proliferation was mediated via MHC class I recognition. Furthermore, the reduced in vivo half-life of IL-15 complex in $\beta 2m^{-/-}$ mice may explain the decreased response of CD44^{hi} polyclonal CD8 T cells to IL-15 complex in these mice.

TCR avidity for self-MHC controls responsiveness to IL-15 complex

Not all naive CD8 T cells respond equally well to a lymphopenic environment (4), and this finding has been linked to TCR avidity (28). Because the CD8 T cell response to IL-15 complex was also MHC class I dependent, it was of interest to test whether relative TCR avidities affected the response. We first examined whether the response of naive CD8 T cells to IL-15/IL-15Rα complex treatment exhibited heterogeneity. CFSE-labeled CD45.1 CD44^{lo} naive polyclonal B6 CD8 T cells were transferred into normal CD45.2 B6 hosts on day -1 that were then treated with PBS or IL-15 complex on day 0. Concurrently, CFSE-labeled CD45.1 CD44^{lo} naive polyclonal CD8 T cells were also transferred into CD45.2 B6 hosts sublethally irradiated (550 rad) 1 d earlier. At 4-5 d posttreatment, splenocytes were examined for CFSE dilution. A robust proliferative response was seen only when the mice were exposed to IL-15 complex (Fig. 3A). Interestingly, the polyclonal CD8 T cell population divided in an asynchronous fashion, in contrast to OT-I cells (Fig. 1), with some cells undergoing extensive division and others barely dividing. Thus, not all naive endogenous CD8 T cells responded to IL-15 complex to the same extent. Similarly, although the proliferative response was not as robust as in IL-15 complex-treated hosts, polyclonal CD8 T cells also exhibited an asynchronous division in sublethally irradiated hosts (Fig. 3A).

To more closely examine how TCR avidity may influence the response to IL-15 complex, we compared the response of naive TCR transgenic CD8 T cells of different specificities and distinct TCR avidities (28). To this end, naive CFSE-labeled CD44^{lo} CD45.1 OT-I (specific for SIINFEKL peptide presented by H-2K^b), F5 (specific for influenza virus NP peptide presented by H-2D^b), and C4 (specific for influenza virus HA peptide presented by H-2K^b) TCR transgenic CD8 T cells were transferred into CD45.2 B6 or B10.D2 (for C4) mice on day -1. On day 0, B6 hosts were treated with PBS or IL-15 complex. TCR transgenic T cells were also transferred into CD45.2 B6 mice sublethally irradiated (550 rad) 1 d in advance. On day 5 posttreatment, donor cells were analyzed for CFSE dilution. In an immunodeficient environment (sublethally irradiated hosts), OT-I and C4 cells divided extensively, whereas F5 cells did not divide (Fig. 3B). Thus, the TCR avidity of F5 CD8 T cells is insufficient to drive homeostatic proliferation. Interestingly, the response of OT-I, F5, and C4 TCR transgenic CD8 T cells to IL-15 complex paralleled the response to lymphopenia. This was not due to differences in CD122 or CD132 expression levels between the TCR transgenic T cells (data not shown). Compared with lymphopenia-driven expansion, proliferation was better with IL-15 complex treatment in all cases, with some limited proliferation observed with F5 T cells (Fig. 3B).



FIGURE 3. Naive CD8 T cell response to IL-15 complex is heterogeneous. CD44¹⁰ polyclonal CD8 T cells (*A*) or OT-I, F5, or C4 CFSE-labeled CD45.1 CD44¹⁰ CD8 T cells (*B*) were injected i.v. into irradiated (500 rad) or normal CD45.2 B6 or B10.d2 (for C4) mice. One day later, the normal B6 mice received PBS or 2.5 μ g hIL-15 + 15 μ g mouse IL-15R α -Fc i.p. Five days after treatment, donor cell proliferation was examined in the spleen.

IL-15 complex enhances the naive CD8 T cell response to cognate peptide

Providing naive CD8 T cells their cognate peptide during acute homeostatic proliferation enhances their proliferative response (8, 10). Therefore, we examined whether IL-15 complex treatment, along with cognate peptide, would further enhance the naive CD8 T cell response to IL-15 complex. CFSE-labeled CD45.1 CD44^{lo}

6861

OT-I and F5 cells were transferred into separate CD45.2 B6 hosts on day -1. On day 0, hosts were treated with PBS, cognate peptide, IL-15/IL-15Ra, or cognate peptide and IL-15/IL-15Ra. On day 5, donor cell division was measured by flow cytometry. Peptide alone induced significant proliferation in F5 cells, whereas IL-15 complex was a poor inducer of cell division. However, the F5 cell response was greatly enhanced when IL-15 complex and cognate peptide were administered (Fig. 4A). Similar results were obtained with OT-I cells, although their response to IL-15 complex alone was much greater than that of F5 cells (Fig. 4B). We also examined the accumulation of the different cell populations in each case. Peptide treatment alone, although inducing cell division, did not result in significant accumulation of responding cells, likely due to deletion (Fig. 4). Treatment with the combination of cognate peptide and IL-15 complex seemed to be synergistic, resulting in substantial accumulation of cells. In additional studies in which OT-I cells and F5 cells were cotransferred, OT-I cells outcompeted F5 in each of the responses (Supplemental Fig. 1). This finding further supported the concept that TCR avidity was controlling the response to IL-15/IL-15R α complex. Nevertheless, these results suggested that IL-15 complex acts selectively on Ag-activated CD8 T cells and could serve as an adjuvant for Ag-specific responses. Furthermore, the cellaccumulation data illustrated that naive CD8 T cells normally unresponsive to IL-15 complex were more responsive when TCR was engaged.

We also examined the effect of peptide and IL-15 complex treatment on T cell migration to nonlymphoid tissues. CD45.1 CD44^{lo} naive OT-I cells were transferred into CD45.2 B6 hosts on day -1, and on day 0 these mice were treated with PBS, SIIN-FEKL peptide, IL-15 complex, or SIINFEKL peptide and IL-15 complex (peptide/IL-15 complex). Five days posttreatment, mice were sacrificed, and various tissues were examined for the presence of the OT-I donor population. IL-15 complex treatment increased OT-I cell numbers in the lung, as well as in the spleen. An

FIGURE 4. TCR avidity controls the CD8 T cell response to IL-15 complex. F5 (*A*) or OT-I CFSE-labeled CD45.1 CD44^{lo} (*B*) CD8 T cells were injected i.v. into B6 mice. One day later, mice were treated i.p. with PBS, IL-15 complex, or with F5-NP peptide (100 μ g) or SIINFEKL (6.25 μ g), or with peptide plus IL-15 complex. Five days after treatment, donor cells were examined for proliferation and total cell numbers in the spleen. Data are representative of two similar experiments (*n* = 3).



FIGURE 5. Cognate peptide increases and alters the naive CD8 T cell response to IL-15 complex. A, CD45.1 CD4410 OT-I CD8 T cells were i.v. transferred into CD45.2 B6 mice. One day later, mice were treated i.p. with PBS, 100 µg SIINFEKL peptide, 2.5 µg hIL-15 + 15 μg IL-15Rα-Fc, or 100 μg SIINFEKL and 2.5 µg IL-15 + 15 µg IL-15Rα-Fc. Donor cell accumulation was examined in the spleen, lungs, and brain. B. Donor cells from the spleen were analyzed for expression of the indicated proteins by flow cytometry. Shaded graph = PBS, thin line = IL-15 + IL-15R α -Fc, thick line = peptide, dashed line = peptide + IL-15 + IL-15R α -Fc. Data are representative of at least two experiments (n = 3-4).



even larger OT-I donor population was present in both tissues of mice treated with peptide/IL-15 complex compared with IL-15 complex treatment alone (Fig. 5). A similar pattern was seen in the liver (data not shown). Comparable results were obtained when F5 cells were transferred and the treatments administered, although, as expected, fewer total F5 cells compared with OT-I cells, were recovered from the organs (Supplemental Fig. 2). When the brain, a more immunoprivileged tissue (37, 38), was examined, very few OT-I CD8 T cells were present in control, peptide- or IL-15 complex-treated mice. In contrast, a sizable population of CD8 T cells was present in the brains of peptide/IL-15 complex-treated mice (Fig. 5).

The previous data suggested that IL-15 complex treatment, along with peptide immunization, enhanced lymphocyte migration. Therefore, we examined the expression of various adhesion molecules after treatment. All treatments uniformly upregulated CD11a (Fig. 5*B*). Treatment with IL-15 complex alone weakly upregulated CD43 and did not induce CD62L downregulation, but it modestly enhanced CD44 expression. Peptide treatment alone strongly induced CD43 expression, modestly downregulated CD62L expression, and greatly upregulated CD44 levels. Peptide/IL-15 complex was the most effective at CD43 induction and CD62L downregulation, and it strongly induced CD44 expression. Thus, IL-15 complex enhanced the proliferative response to cognate Ag, and the combination of IL-15 complex and Ag also resulted in the modulation of adhesion receptors involved in migration and enhanced function.

We performed serial adoptive transfers to formally test whether IL-15 complex altered the migration potential of the responding CD8 T cells. CD45.1-naive CD44^{lo} OT-I cells were transferred into CD45.2 B6 hosts on day -1 and then treated with IL-15 complex,

peptide/IL-15 complex, or VSV-OVA on day 0. VSV-OVA infection causes similar changes in cell-surface phenotype as does peptide/IL-15 complex (data not shown) and was used as a positive control for migration potential to tertiary sites. On day 4 posttreatment, splenocytes were harvested and enriched for donor OT-I cells. A total of 1×10^{6} CD45.1 OT-I cells from each group, including naive CD44^{lo} OT-I cells as a control, were transferred into secondary CD45.2 B6 hosts. Two days after secondary transfer, hosts were sacrificed, and the presence of donor OT-I cells was determined in several tissues (Fig. 6). Naive OT-I cells were mainly found in the spleen and pLNs. In contrast, VSV-OVA-experienced OT-I cells were found in the spleen, as well as in the lungs and liver, with few present in the pLNs. Cells from IL-15 complex-treated mice behaved similarly to naive CD8 T cells, whereas the migration pattern of peptide/IL-15 complextreated cells resembled that of virus-activated OT-I cells. Overall, the data support the notion that IL-15 complex treatment can augment CD8 T cell activation and enhance migratory capabilities, both of which are necessary outcomes of effective immunotherapy.

Discussion

Our previous work highlighted the intriguing ability of IL-15R α complexed IL-15, but not IL-15 alone, to activate naive CD8 T cells (16). This response was Ag independent, but we now show that the process required expression of MHC class I. Furthermore, the ability of a naive CD8 T cell to respond to IL-15 complex correlated with TCR avidity for self-peptide:MHC class I. Intriguingly, the response was augmented, as well as altered, by providing a stronger TCR stimulus. In particular, IL-15 complex

FIGURE 6. Activation with cognate peptide and IL-15 complex alters the migratory ability of CD8 T cells. A total of 6×10^6 congenic naive OT-I cells were transferred into B6 hosts on day -1 and then were treated i.p. with 2.5 μ g IL-15 + 15 μ g IL-15R α -Fc, 100 µg SIINFEKL and 2.5 µg IL-15 + 15 µg IL-15R α -Fc, or 1 \times 10⁵ PFU VSV-OVA on day 0. Four days posttreatment, splenocytes were harvested and enriched for donor OT-I cells. OT-I cells (1×10^6) from each group, including fresh CD44^{lo} naive OT-I cells, were transferred i.v. into secondary untreated B6 hosts. Two days after secondary transfer, mice were sacrificed, and the presence of donor OT-I cells was determined in spleen, pLNs, liver, and lung. Data are depicted as the percentage of cells in a tissue out of four tissues tested and are representative of two experiments (n = 3).



induced a lymphoid-migration profile, whereas agonist peptide/IL-15 complex induced a nonlymphoid-migration profile. These latter results point out some important considerations for the use of IL-15 complex as a therapeutic and may help to explain some recent findings.

Recent work revealed similarities in the naive CD8 T cell response to IL-15 complex and IL- 2α /IL-2 complex (IL-2 complex). Similar to the effects of IL-15 complex treatment (16), naive CD8 T cells treated with IL-2 complex in vivo proliferate, acquire an activated/memory phenotype, produce effector cytokines, and kill in an Ag-specific manner (39). In apparent contrast to our findings, IL-2 complex-driven naive CD8 T cell activation is classical MHC class I independent (39). However, this study used H-2K^b and H-2D^b double-deficient mice, which still express other MHC class I molecules. Indeed, another report found that the CD8 T cell response to IL-2 complex is MHC class I dependent when using $K^{b^{-}/-}$ $D^{b^{-}/-}$ $\beta 2m^{-/-}$ triple-knockout mice (35), ensuring that classical and nonclassical MHC class I molecules are absent. An additional study found that in CD132^{-/-} ($\gamma C^{-/-}$) mice, in which IL-2 and IL-15 levels are abnormally high, naive CD8 T cell proliferation is only modestly affected by the absence of MHC class I (40), perhaps as a result of the combined effects of both cytokines. Overall, these findings further illustrate the close relationship between factors controlling lymphopenia-driven and cytokine complex-driven naive CD8 T cell activation.

One caveat with the use of $\beta 2m^{-/-}$ mice to study the response to an IgG-Fc region-containing molecule is the potential role of FcRn in IgG catabolism (36). This issue could be a confounding factor in our studies and that of Cho et al. (35). In fact, we found that the in vivo half-life of IL-15 was modestly and similarly reduced in $\beta 2m^{-/-}$ and FcRn^{-/-} mice. However, IL-15 complexdriven naive CD8 T cell proliferation remained robust in FcRn^{-/-} mice, whereas it was substantially diminished in $\beta 2m^{-/-}$ mice. Thus, the absence of FcRn cannot account for the lack of proliferation seen in $\beta 2m^{-/-}$ mice. Not surprisingly, the activity of IL-2/ α -IL-2 was also recently shown to be dependent, in part, on FcRn (41).

Our data indicated that not all naive CD8 T cells responded with the same intensity to IL-15 complex. A similar spectrum of proliferation is seen when naive CD8 T cells are placed in a lymphopenic environment (9). Recent work noted that TCR avidity for self-peptide:MHC class I regulates the CD8 T cell response to lymphopenia. Thus, in a lymphopenic environment, T cells with high-avidity TCRs proliferate more vigorously than do T cells with lower-affinity TCRs (28). Our results demonstrated a direct correlation between the relative ability of a naive CD8 T cell to respond to IL-15 complex or to lymphopenia. Furthermore, the response to IL-15 complex positively correlated with TCR avidity, as measured by acute homeostatic proliferation. Greater TCR signaling may confer heightened responsiveness to homeostatic cytokines, such as IL-7, especially in more competitive situations (28, 42). Thus, the cytokine-driven response to lymphopenia also requires a level of tonic TCR signaling, although there is some evidence of TCR-independent cytokine-mediated proliferation in lymphopenic settings (29). There is also a link between IL-15 adjuvanticity and TCR avidity, because IL-15 administration causes the outgrowth of higher-avidity CTLs in response to infection (43). Our cotransfer studies reiterated the concept that high-avidity clones may outpace low-avidity clones in response to IL-15 complex in the presence or absence of Ag. Given these findings, IL-15 complex may operate as an effective immunotherapy by preferentially stimulating the expansion of high-avidity naive or effector CD8 T cells, in part through TCR- and MHCmediated signals.

Of particular interest was our finding that IL-15 complex and peptide/IL-15 complex treatments caused distinct changes in the expression of surface markers associated with migration. Although complex and peptide/IL-15 complex caused upregulation of CD11a and CD44, only peptide/IL-15 complex caused significant downregulation of CD62L and upregulation of CD43. CD43 plays a role in the migration of effector cells from secondary lymphoid tissues to tertiary sites of inflammation, in particular the brain (44, 45). Furthermore, the concomitant downregulation of CD62L and upregulation of CD44 also promote migration to tertiary sites (46-48). Overall, the changes in surface expression of the receptors correlated with the migratory behavior of the cells. CD8 T cells activated by peptide/IL-15 complex treatment migrated more readily to nonlymphoid tissues compared with cells activated with IL-15 complex alone. The efficacy of IL-15 complex in tumor immunotherapy may hinge, in some cases, on the ability of IL-15 complex to alter effector cell migration. For example, partial regression of solid tumors is induced by IL-15 complex by

activation of intratumor tumor-specific CD8 T cells (19). The investigators noted that IL-15 complex treatment more effectively stimulated intratumor CD8 T cells than peripheral CD8 T cells, and IL-15 complex treatment did not enhance CD8 T cell migration into tumors. Given our current results, the robust response of intratumor CD8 T cells may be explained by access and interaction of intratumor, but not peripheral, CD8 T cells with cognate Ag. Without access to tumor Ag, IL-15 complex-activated tumor-specific CD8 T cells may not efficiently enter pre-existing tumors because of the lack of effective modulation of homing receptors that promote extralymphoid migration. Thus, stronger TCR stimulation, in conjunction with IL-15R signaling, may help to spur migration of CTLs into tertiary tissues and, possibly, tumors.

Overall, our findings highlight the ability of IL-15 complex to specifically induce activation and proliferation of naive CD8 T cells expressing high-avidity TCRs. This process is dependent on the expression of self-peptide:MHC class I. This feature of the IL-15 complex response may mimic the action of IL-15 during immune responses, in which higher-avidity T cells may compete more effectively for homeostatic cytokines. IL-15 complex holds promise as an immunotherapy, but further study of the synergy between TCR and IL-15R signaling is required to aid in the design of therapeutic protocols.

Disclosures

T.A.S. and L.L. have a patent pending related to the IL-15 complex and have received royalties from Marine Polymers Technology, Inc. All other authors have no financial conflicts of interest.

References

- Nesic, D., and S. Vukmanovic. 1998. MHC class I is required for peripheral accumulation of CD8+ thymic emigrants. J. Immunol. 160: 3705–3712.
- Murali-Krishna, K., L. L. Lau, S. Sambhara, F. Lemonnier, J. Altman, and R. Ahmed. 1999. Persistence of memory CD8 T cells in MHC class Ideficient mice. *Science* 286: 1377–1381.
- Tan, J. T., B. Ernst, W. C. Kieper, E. LeRoy, J. Sprent, and C. D. Surh. 2002. Interleukin (IL)-15 and IL-7 jointly regulate homeostatic proliferation of memory phenotype CD8+ cells but are not required for memory phenotype CD4+ cells. J. Exp. Med. 195: 1523–1532.
- Schluns, K. S., W. C. Kieper, S. C. Jameson, and L. Lefrançois. 2000. Interleukin-7 mediates the homeostasis of naïve and memory CD8 T cells in vivo. *Nat. Immunol.* 1: 426–432.
- Goldrath, A. W., P. V. Sivakumar, M. Glaccum, M. K. Kennedy, M. J. Bevan, C. Benoist, D. Mathis, and E. A. Butz. 2002. Cytokine requirements for acute and Basal homeostatic proliferation of naive and memory CD8+ T cells. *J. Exp. Med.* 195: 1515–1522.
- Schluns, K. S., K. D. Klonowski, and L. Lefrançois. 2004. Transregulation of memory CD8 T-cell proliferation by IL-15Ralpha+ bone marrow-derived cells. *Blood* 103: 988–994.
- Wu, T. S., J. M. Lee, Y. G. Lai, J. C. Hsu, C. Y. Tsai, Y. H. Lee, and N. S. Liao. 2002. Reduced expression of Bcl-2 in CD8+ T cells deficient in the IL-15 receptor alpha-chain. *J. Immunol.* 168: 705–712.
- Goldrath, A. W., and M. J. Bevan. 1999. Low-affinity ligands for the TCR drive proliferation of mature CD8+ T cells in lymphopenic hosts. *Immunity* 11: 183– 190.
- Ernst, B., D. S. Lee, J. M. Chang, J. Sprent, and C. D. Surh. 1999. The peptide ligands mediating positive selection in the thymus control T cell survival and homeostatic proliferation in the periphery. *Immunity* 11: 173–181.
- Kieper, W. C., and S. C. Jameson. 1999. Homeostatic expansion and phenotypic conversion of naïve T cells in response to self peptide/MHC ligands. *Proc. Natl. Acad. Sci. USA* 96: 13306–13311.
- Tan, J. T., E. Dudl, E. LeRoy, R. Murray, J. Sprent, K. I. Weinberg, and C. D. Surh. 2001. IL-7 is critical for homeostatic proliferation and survival of naive T cells. *Proc. Natl. Acad. Sci. USA* 98: 8732–8737.
- Sandau, M. M., C. J. Winstead, and S. C. Jameson. 2007. IL-15 is required for sustained lymphopenia-driven proliferation and accumulation of CD8 T cells. J. Immunol. 179: 120–125.
- Dubois, S., J. Mariner, T. A. Waldmann, and Y. Tagaya. 2002. IL-15Ralpha recycles and presents IL-15 In trans to neighboring cells. *Immunity* 17: 537–547.
- Burkett, P. R., R. Koka, M. Chien, S. Chai, D. L. Boone, and A. Ma. 2004. Coordinate expression and trans presentation of interleukin (IL)-15Ralpha and IL-15 supports natural killer cell and memory CD8+ T cell homeostasis. *J. Exp. Med.* 200: 825–834.
- Mortier, E., A. Quéméner, P. Vusio, I. Lorenzen, Y. Boublik, J. Grötzinger, A. Plet, and Y. Jacques. 2006. Soluble interleukin-15 receptor alpha (IL-15R

alpha)-sushi as a selective and potent agonist of IL-15 action through IL-15R beta/gamma. Hyperagonist IL-15 x IL-15R alpha fusion proteins. *J. Biol. Chem.* 281: 1612–1619.

- Stoklasek, T. A., K. S. Schluns, and L. Lefrançois. 2006. Combined IL-15/IL-15Ralpha immunotherapy maximizes IL-15 activity in vivo. J. Immunol. 177: 6072–6080.
- Rubinstein, M. P., M. Kovar, J. F. Purton, J. H. Cho, O. Boyman, C. D. Surh, and J. Sprent. 2006. Converting IL-15 to a superagonist by binding to soluble IL-15Rα. Proc. Natl. Acad. Sci. USA 103: 9166–9171.
- Dubois, S., H. J. Patel, M. Zhang, T. A. Waldmann, and J. R. Müller. 2008. Preassociation of IL-15 with IL-15R alpha-IgG1-Fc enhances its activity on proliferation of NK and CD8+/CD44high T cells and its antitumor action. J. Immunol. 180: 2099–2106.
- Epardaud, M., K. G. Elpek, M. P. Rubinstein, A. R. Yonekura, A. Bellemare-Pelletier, R. Bronson, J. A. Hamerman, A. W. Goldrath, and S. J. Turley. 2008. Interleukin-15/interleukin-15R alpha complexes promote destruction of established tumors by reviving tumor-resident CD8+ T cells. *Cancer Res.* 68: 2972– 2983.
- Goldrath, A. W., L. Y. Bogatzki, and M. J. Bevan. 2000. Naive T cells transiently acquire a memory-like phenotype during homeostasis-driven proliferation. J. Exp. Med. 192: 557–564.
- Murali-Krishna, K., and R. Ahmed. 2000. Cutting edge: naive T cells masquerading as memory cells. J. Immunol. 165: 1733–1737.
- Becker, T. C., E. J. Wherry, D. Boone, K. Murali-Krishna, R. Antia, A. Ma, and R. Ahmed. 2002. Interleukin 15 is required for proliferative renewal of virusspecific memory CD8 T cells. J. Exp. Med. 195: 1541–1548.
- Cho, B. K., V. P. Rao, Q. Ge, H. N. Eisen, and J. Chen. 2000. Homeostasisstimulated proliferation drives naive T cells to differentiate directly into memory T cells. J. Exp. Med. 192: 549–556.
- Ge, Q., H. Hu, H. N. Eisen, and J. Chen. 2002. Different contributions of thymopoiesis and homeostasis-driven proliferation to the reconstitution of naive and memory T cell compartments. *Proc. Natl. Acad. Sci. USA* 99: 2989– 2994.
- Brossart, P., A. W. Goldrath, E. A. Butz, S. Martin, and M. J. Bevan. 1997. Virusmediated delivery of antigenic epitopes into dendritic cells as a means to induce CTL. J. Immunol. 158: 3270–3276.
- Kieper, W. C., A. Troy, J. T. Burghardt, C. Ramsey, J. Y. Lee, H. Q. Jiang, W. Dummer, H. Shen, J. J. Cebra, and C. D. Surh. 2005. Recent immune status determines the source of antigens that drive homeostatic T cell expansion. *J. Immunol.* 174: 3158–3163.
- Kassiotis, G., R. Zamoyska, and B. Stockinger. 2003. Involvement of avidity for major histocompatibility complex in homeostasis of naive and memory T cells. *J. Exp. Med.* 197: 1007–1016.
- 28. Kieper, W. C., J. T. Burghardt, and C. D. Surh. 2004. A role for TCR affinity in regulating naive T cell homeostasis. J. Immunol. 172: 40–44.
- Hao, Y., N. Legrand, and A. A. Freitas. 2006. The clone size of peripheral CD8 T cells is regulated by TCR promiscuity. J. Exp. Med. 203: 1643– 1649.
- Roopenian, D. C., G. J. Christianson, T. J. Sproule, A. C. Brown, S. Akilesh, N. Jung, S. Petkova, L. Avanessian, E. Y. Choi, D. J. Shaffer, et al. 2003. The MHC class I-like IgG receptor controls perinatal IgG transport, IgG homeostasis, and fate of IgG-Fc-coupled drugs. J. Immunol. 170: 3528–3533.
- Moskophidis, D., and D. Kioussis. 1998. Contribution of virus-specific CD8+ cytotoxic T cells to virus clearance or pathologic manifestations of influenza virus infection in a T cell receptor transgenic mouse model. *J. Exp. Med.* 188: 223–232.
- Morgan, D. J., R. Liblau, B. Scott, S. Fleck, H. O. McDevitt, N. Sarvetnick, D. Lo, and L. A. Sherman. 1996. CD8(+) T cell-mediated spontaneous diabetes in neonatal mice. J. Immunol. 157: 978–983.
- Hamilton, S. E., M. C. Wolkers, S. P. Schoenberger, and S. C. Jameson. 2006. The generation of protective memory-like CD8+ T cells during homeostatic proliferation requires CD4+ T cells. *Nat. Immunol.* 7: 475–481.
- Lyons, A. B., and C. R. Parish. 1994. Determination of lymphocyte division by flow cytometry. J. Immunol. Methods 171: 131–137.
- Cho, J. H., O. Boyman, H. O. Kim, B. Hahm, M. P. Rubinstein, C. Ramsey, D. M. Kim, C. D. Surh, and J. Sprent. 2007. An intense form of homeostatic proliferation of naive CD8+ cells driven by IL-2. *J. Exp. Med.* 204: 1787– 1801.
- Roopenian, D. C., and S. Akilesh. 2007. FcRn: the neonatal Fc receptor comes of age. Nat. Rev. Immunol. 7: 715–725.
- Klonowski, K. D., K. J. Williams, A. L. Marzo, D. A. Blair, E. G. Lingenheld, and L. Lefrançois. 2004. Dynamics of blood-borne CD8 memory T cell migration in vivo. *Immunity* 20: 551–562.
- Mrass, P., and W. Weninger. 2006. Immune cell migration as a means to control immune privilege: lessons from the CNS and tumors. *Immunol. Rev.* 213: 195– 212.
- Kamimura, D., and M. J. Bevan. 2007. Naive CD8+ T cells differentiate into protective memory-like cells after IL-2 anti IL-2 complex treatment in vivo. J. Exp. Med. 204: 1803–1812.
- Ramsey, C., M. P. Rubinstein, D. M. Kim, J. H. Cho, J. Sprent, and C. D. Surh. 2008. The lymphopenic environment of CD132 (common gamma-chain)deficient hosts elicits rapid homeostatic proliferation of naive T cells via IL-15. J. Immunol. 180: 5320–5326.
- 41. Létourneau, S., E. M. van Leeuwen, C. Krieg, C. Martin, G. Pantaleo, J. Sprent, C. D. Surh, and O. Boyman. 2010. IL-2/anti-IL-2 antibody complexes show strong biological activity by avoiding interaction with IL-2 receptor alpha subunit CD25. *Proc. Natl. Acad. Sci. USA* 107: 2171–2176.

- Seddon, B., and R. Zamoyska. 2002. TCR and IL-7 receptor signals can operate independently or synergize to promote lymphopenia-induced expansion of naive T cells. *J. Immunol.* 169: 3752–3759.
- Oh, S., L. P. Perera, D. S. Burke, T. A. Waldmann, and J. A. Berzofsky. 2004. IL-15/IL-15Ralpha-mediated avidity maturation of memory CD8+ T cells. *Proc. Natl. Acad. Sci. USA* 101: 15154–15159.
- 44. Stockton, B. M., G. Cheng, N. Manjunath, B. Ardman, and U. H. von Andrian. 1998. Negative regulation of T cell homing by CD43. *Immunity* 8: 373–381.
- Onami, T. M., L. E. Harrington, M. A. Williams, M. Galvan, C. P. Larsen, T. C. Pearson, N. Manjunath, L. G. Baum, B. D. Pearce, and R. Ahmed. 2002.

Dynamic regulation of T cell immunity by CD43. J. Immunol. 168: 6022-6031.

- Arbonés, M. L., D. C. Ord, K. Ley, H. Ratech, C. Maynard-Curry, G. Otten, D. J. Capon, and T. F. Tedder. 1994. Lymphocyte homing and leukocyte rolling and migration are impaired in L-selectin-deficient mice. *Immunity* 1: 247–260.
- DeGrendele, H. C., P. Estess, and M. H. Siegelman. 1997. Requirement for CD44 in activated T cell extravasation into an inflammatory site. *Science* 278: 672–675.
- Stoop, R., I. Gál, T. T. Glant, J. D. McNeish, and K. Mikecz. 2002. Trafficking of CD44-deficient murine lymphocytes under normal and inflammatory conditions. *Eur. J. Immunol.* 32: 2532–2542.