

Product(s)	Administration**	Storage/Stability	
Humate-P	Humate P given over 3-20 min (Max 4mL/min)	Prior to reconstitution: 2-8°C or room temp (≤25C), do not freeze.	Reconstitution: Bring to room temperature and reconstitute with provided diluent. Swirl, do not shake or filter. After reconstitution: Room temp (≤25°C), use within 3hr
FEIBA NH	FEIBA max rate of 2 units/kg/min		
BeneFix	BeneFix given at a rate of 2-4mL/min over several minutes	Prior to reconstitution: 2-8°C, do not freeze. May be stored at room temp (≤25C) for up to 6 months	
Kogenate	Over 1 to 15 minutes IVP unless ordered as Continuous Infusion	Prior to reconstitution: 2-8°C, do not freeze, Helixate FS may be stored at room temp ≤ 25C, up to 12 months.	After reconstitution: Room temp (≤25°C), use within 3hr Continuous infusion of Helixate FS or Recombinate 10 IU/mL dilution in NS. Dose based on replacement requirements. Infuse within 12hrs
NovoSeven RT	given as IVP over 2-5 minutes (can also be ordered as Continuous Infusion)	Prior to reconstitution: <25°C	After reconstitution: Room temp (≤25°C) of refrigerated IVP: Use within 3hr; do not freeze; do not store in syringes CI: Do not dilute. Infuse within 12 hours (stability questionable beyond 12-24 hours)

**Refer to JDH/UConn IV Guidelines for further information. Do not use an inline filter. Flush with Normal Saline, Store all products in original packaging

Humate-P Dosing for VWD Hemorrhage DOSED IN VWF:RCO ONLY		
VWD Type	Severity	Dosage (IU VWF:RCO/kg)
Type 1 – Mild (VWF:RCO >30%)	Minor	Use DDAVP IV 0.3mcg/kg by slow infusion
	Minor w/o DDAVP or Major	Load 40-60 IU/kg then 40-50IU/kg q8-12hr x3 days ¹ , then 40-50IU/kg qday ²
Type 1 – Moderate/ Severe (VWF:RCO <30%)	Minor	40-50 IU/kg (x1-2 doses)
	Major	Load 50-75 IU/kg then 40-60 IU/kg q8-12hr x3 days ¹ , then 40-60 IU/kg qday ²
Type 2 and 3 (all)	Minor	40-50 IU/kg (x1-2 doses)
	Major	Load 60-80 IU/kg then 40-60 IU/kg q8-12hr x3 days ¹ , then 40-60 IU/kg qday ²

¹To maintain VWF:RCO trough level >50%

²For up to 7 days

References: LexiComp, NovoSeven RT (NovoNordisk) package insert <http://www.novo-pi.com/novosevenrt.pdf>
Recombine (Baxter) package insert <http://factorviiia.com/pi.pdf>
BeneFix (Wyeth) package insert <http://www.pfizerpro.com/resources/minisites/hemophilia/docs/BeneFIX-PI.pdf>
FEIBA (Baxter) package insert http://www.feiba.com/us/forms/feiba_nf_pi.pdf
Humate-P (CSLBehring) package insert <http://www.humate-p.com/Professional/Prescribing-Information.aspx>
Thigpen and Limdi, Reversal of Oral Anticoagulation. *Pharmacotherapy*, vol 33, issue 11, pp. 1199-1213, November, 2013
KCentra package insert, <http://www.kcentra.com/Kcentra-resources.aspx>

BILLING INFO

- EVERY bolus and IV factor dispensed including Kcentra **must be entered in the Cubixx log clearly and neatly and initialed clearly by the dispensing Pharmacist.**
- Entering pharmacists must physically enter in the number of labels needed to ensure doses are sent, using ... labels for inpatients.
- Billing personnel will verify all factor billing with the log, and MAK or the MAR.
- At present, Factor Products are done in Dose Edge
- Bolus doses in excess of 3 vials should be drawn up and labeled by Pharmacy once need of dose is verified.

UConn HEALTH

UConn JOHN DEMPSEY
HOSPITAL

Department of Pharmacy, November 2018
Revised by Ruth Kalish, RPH

UConn Health
Department of Pharmacy
**Procoagulant
(Factor)
Agent**



Helpful Hints

For Factors VIII and IX, IU/kg dose is a general guideline, as dosage should be based on desired/expected goal factor concentration.

For Factor VIII, 1 IU/kg can be expected to raise the factor levels by 2 percentage points. For Factor IX, 1 IU/kg can be expected to raise factor levels by 0.75 percentage points

Humate P is the only product available for patients with von Willebrand Disease (VWD). Dose is based on VWF:RCO units only for VWD. VWF units varies in ratio to Factor VIII units with an average of 2.4:1.

* Per Pharmacy & Therapeutics committee, Pharmacy can adjust dose within +/- 10% based on current dosage vial sizes in pharmacy to prevent waste. Rounding up to the nearest vial size is preferred.

ASD Healthcare has provided a Cubixx refrigerator to store the consignment product in. If there is a mechanical issue with the refrigerator contact card is taped to the refrigerator.

The pharmacy will be billed for the product when it has been removed from the Cubixx unit for more than 180 minutes.

When the inventory falls below the designated par levels, the Cubixx unit will automatically generate an order. The turnaround time for system generated orders is 2 days.

Do not deface any packaging for all factor agents (e.g. write on or label the box, remove the filter, remove the package insert) or JDH Pharmacy will be required to pay for the product since they are on consignment. Store products in the original package to protect from light.

Pharmacists ONLY: If an order needs to be placed urgently off hours, call 1-800-746-6273. The operator will have an associate call back. Identify yourself as John Dempsey Hospital, acct. # 206267.

UConn
HEALTH
UConn JOHN DEMPSEY
HOSPITAL

Department of Pharmacy

Product	Half-Life (h)	How Supplied	Indications: FDA-Approved (<i>Unapproved</i>)	Type of Bleed	Dose	Interval	Duration	Goal Factor Concentration
BeneFIX * Factor IX (recombinant) Pfizer 1-800-505-4426	11-36	250, 500, 1000, and 2000 IU vials w/ 5ml diluent Final conc: 50, 100, 200, 400IU/mL	Hemophilia B Number of factor IX IU required = patient weight (kg) x desired factor IX level increase (as % normal or IU/dL) x 1.3 (as units/kg or units/dL)	Minor	Up to 40 IU/kg	12-24hr	Until resolved	20-30%
				Moderate	Up to 65 IU/kg	12-24hr	up to 7 days	25-50%
				Major	Up to 130 IU/kg	12-24hr	7-10 days or until resolved	50-100%
				Surgery	Up to 130 IU/kg pre-op, then less	12-24hr PRN to Keep ≤50%	7-14 day or until resolved	Pre-op: 100% Post-op ≤ 50%
FEIBA NF* Activated Prothrombin Complex Concentrate (plasma-derived) Baxter Healthcare (800) 422-9837	4-7 Duration = 8-12h	approx. 500IU or 1000 IU/ 20ml and approx. 2500 IU/50ml vials	Hemophilia A/B with inhibitors (<i>acquired hemophilia with factor VIII or factor IX inhibitor titers >5 Bethesda units (BU) Treatment of life threatening bleeds associated with dabigatran</i>)	Joint	50-100 Feiba units/kg	12hr DIC or thrombosis may occur if 200unit max dose exceeded	Consult specialist if ineffective or >3 doses in 24hr	NA
				Mucous membrane	50-100 Feiba units/kg			
				Soft tissue		12hr	50 unit/kg dose can be q6h	
				Severe (CNS)	100 Feiba units/kg Max of 200 units/kg			
NovoSeven RT Factor VIIa (recombinant) Novo Nordisk (800) 727-6500	1.7-3.1	1mg, 2mg, 5mg, 8mg vials as 1mg/mL	Hemophilia A or B with inhibitors to factor VIII or factor IX, acquired hemophilia, or congenital factor VII deficiency (<i>warfarin-related intracerebral hemorrhage; treatment of refractory bleeding after cardiac surgery in nonhemophiliac patients</i>) CI = 1 mg/ml FVIIa diluent	Hemophilia A or B	90 mcg/kg	q2hr initially. Cont. Post-op until healed Q2h up to 5 days, then 2-6 hrs.	Until hemostasis achieved	NA
				Acquired hemoph.	70-90 mcg/kg			
				Congenital factor VII deficiency	IV Push: 15-30mcg/kg	IVP q4-6hr until hemostasis, then CI based on levels		
					Continuous, undiluted 12-50mcg/kg/hr			
Helixate FS * Factor VIII (recombinant) CSL Behring (800) 504-5434	11-15	250, 500, 1000, 2000 units/ 2.5ml Final conc: 100IU/mL, 200IU/mL, 400IU/mL, 800IU/ml	Hemophilia A (bleeding/surgery) Also prophylaxis of joint bleeding and to reduce risk of joint damage in children with hemophilia A with no pre-existing joint damage Wt (kg) x desired % increase/2 = dose Continuous infusion = 10 units/ml NS JDH does not carry Recombinate*, but dosing guidelines are similar	Minor	10-20 IU/kg	12-24hr	Until resolved	20-40%
				Moderate/minor surg	15-30 IU/kg	12-24hr	3 days or until-resolved	30-60%
				Major	40-50 IU/kg f/b 20-25IU/kg	8-12hr	3-7 days or until bleed resolved	80-100%
				Major Surgery	50 IU/kg load CI dose based on levels	6-24hr	10-14 days or until healed	Pre-op: 100% Monitor levels
Humate-P * Factor VIII (plasma-derived) CSL Behring (800) 504-5434 See Guidelines for Von Willebrand's Ds on reverse	8-28 (HA)	250/600IU/5ml 500/1200IU/10 ml 1000/2400IU/15 ml	Hemophilia A (classical hemophilia) (this table); Not preferred agent. See Reverse for Severe VWD , including mild or moderate disease where use of desmopressin is known or suspected to be inadequate Note: <i>Not indicated for the prophylaxis of spontaneous bleeding episodes.</i>	Minor	15 IU FVIII/kg f/b 7.5 IU FVIII/kg	12-24hr	Until resolved	30%
				Moderate	25 IU FVIII/kg f/b 15 IU FVIII/kg	8-24hr	3-7 days or until resolved	30-50%
	3-34 (VWD)	VIII/VWF:RCo Ratio Approx 2.4:1		Surgery or Major	40-50 IU FVIII/kg f/b 20-25 IU FVIII/kg	8-24hr	7-14days or until resolved	Pre-op + initial Post-op: 80-100% x7days f/b 30-50% x7d
KCentra* Prothrombin complex concentrate, heat treated, (Factors II, VII, IX, X)	Indication		PreTx INR	Dose (in Factor IX content)	Administration	Caution	Storage	Monitoring
	Urgent reversal of Vit K antagonist induced coagulation factor deficiency in patient with acute major bleed or urgent invasive procedure/surgery. Approx. 500 Units Factor IX activity per vial		2 - <4	25 units/kg max 2500 units	0.12ml/kg/min or approx. 3 units/kg/min. NTE 8.4ml/min	Do not let blood enter syringe to avoid clot formation. Monitor for clotting	Refrigerate. Use within 4 hrs of reconstitution if refrigerated	INR declines within minutes, duration 6-8hrs May affect aPTT
			4 - 6	35 units/kg max 3500 units				
> 6			50 units/kg max 5000 units Do not exceed max dose					