

70 y/o F presents with headache

Samantha Huq, MD, MPH

Leo Wolansky, MD








?

Subarachnoid hemorrhage



Axial NECT shows small foci of hemorrhage in the sulci of the right cerebral convexity



Coronal NECT
confirms
subarachnoid
hemorrhage's
curvilinear
configuration

Subarachnoid hemorrhage (SAH)

Most common cause?

Subarachnoid hemorrhage (SAH)

Most common cause?

Trauma (trick question)

- typically associated with hemorrhagic cortical contusions

Subarachnoid hemorrhage (SAH)

Most common atraumatic (spontaneous)
cause?

Subarachnoid hemorrhage (SAH)

Most common atraumatic (spontaneous) cause?

- Ruptured saccular (berry) aneurysm >85%

Subarachnoid hemorrhage (SAH)

Most common atraumatic (spontaneous) cause?

- Ruptured saccular (berry) aneurysm >85%
- AVM
- dural arteriovenous fistula
- isolated perimesencephalic hemorrhage (unknown cause, probably venous)

Subarachnoid hemorrhage (SAH)

Clinical presentation

- “worst headache of my life”
- Sudden thunderclap headache
- 10% preceded by “sentinel hemorrhage” = self limiting SAH+ headache in preceding days/weeks & can present with infarction

Subarachnoid hemorrhage (SAH)

Imaging:

CT: gold standard, ER work-horse

– sensitivity is influenced by size & time of bleed.

MRI: when optimized is more sensitive

– CSF hyperintensity on FLAIR

– CSF hypointensity on SWI or GRE

– 3D-FLAIR avoids flow artifacts in cisterns

– DSA is the gold standard for diagnosis of aneurysm

CTA, 90-95% positive if aneurysm $\geq 2\text{mm}$

MRA can detect causative aneurysm

Subarachnoid hemorrhage (SAH)

Rx (if aneurysm)

Coil embolization

Micro-neurosurgical clipping – better for some aneurysms

Vasospasm – Ca²⁺ antagonist, angioplasty

Top differential (Pseudo-SAH)

Severe meningitis

Global cerebral edema (Hyperdense o CT)

Supplemental Oxygen (Hyperintense on FLAIR)

Cisternal flow (Hyperintense on 2D-FT FLAIR)

Old SAH (Hypointense on SWI)

Complications

- Rebleeding
- Hydrocephalus (early & late)
- Cerebral infarction from vasospasm (peak ~ 1 week post-bleed)

References

- Rubino S et al: Outpatient follow-up of nonoperative cerebral contusion and traumatic subarachnoid hemorrhage: does repeat head CT alter clinical decision-making? J Neurosurg. 121(4):944-9, 2014
- Servadei F et al: Traumatic Subarachnoid Hemorrhage. World Neurosurg. ePub, 2014
- Quigley MR et al: The clinical significance of isolated traumatic subarachnoid hemorrhage. J Trauma Acute Care Surg. 74(2):581-4, 2013
- <https://emedicine.medscape.com/article/1164341-overview>