
### 78F with decreased O2 sat


Krithika Srikanthan, MD


















# Saddle Embolus







## CTA Findings

- Embolus (filling defect) straddles bifurcation of PA
- If >50% vascular bed reduction
  - right heart strain:
    - RV dilation (RV:LV diameter width >1)
    - straightening or leftward bulging intraventricular septum
    - contrast reflux into IVC, hepatic veins, azygous vein
  - PAH
    - enlargement of the pulmonary trunk (>3cm)
- <15% embolic events result in infarction wedge shaped opacity
- 50% patients with PE have pleural effusion
- 70% will have lower extremity DVT



## Imaging Recommendations

- CTA PA (study of choice)
  - Pregnant patient with ABNORMAL chest radiograph
- V/Q:
  - Contrast allergy
  - Renal failure
  - Pregnant patient with NORMAL chest radiograph



#### Saddle Embolus

- Incidence among patients diagnosed with PE 2.6%
- May be associated with significant hemodynamic instability, but not always. Mortality not necessarily different.
- Rx:
  - Anticoag
  - Thrombolysis for severely symptomatic patients
  - IVC filter
  - Catheter or surgical embolectomy in select high risk people



#### References

- 1. Statdx
- 2. Radiopaedia
- 3. http://www.sciencedirect.com/science/article/pii/S1755001710000643

