


35 y/o M presents with recent leg trauma

Samantha Huq, MD, MPH









?

Tibial plateau fracture



AP view of the knee demonstrates depression of the lateral tibial plateau with minimally displaced fracture fragment



Subcutaneous soft tissue edema as well as knee joint effusion

Tibial plateau fracture

Tibial plateau fracture: originally termed bumper or fender fracture. Only 25% of tibial plateau fractures result from impact with automobile bumpers.

Most common mechanism of injury is axial loading (e.g. falling from a significant height). Soft tissue injuries (e.g. cruciate and collateral ligaments) occur in approximately 10% of patients. Lateral plateau fractures are more common than the medial plateau. If there are medial plateau fractures, it usually occurs in conjunction with lateral tibial plateau fractures.

Imaging: plain films underestimate the severity of injury. Lipohearthrosis is present. CT is helpful in accurately defining the extent of injury. MRI is helpful in assessment of soft tissue injury around the joint.

Schatzker classification is the system for classifying tibial plateau fractures.

References

- McNamara IR et al: Surgical fixation methods for tibial plateau fractures. Cochrane Database Syst Rev. 9:CD009679, 2015
- Burdin G: Arthroscopic management of tibial plateau fractures: surgical technique. Orthop Traumatol Surg Res. 99(1 Suppl):S208-18, 2013
- Persiani P et al: Risk analysis in tibial plateau fractures: association between severity, treatment and clinical outcome. Musculoskelet Surg. 97(2):131-6, 2013