



A large, stylized leaf graphic in a dark blue color, positioned on the left side of the slide. The leaf has a prominent vein structure and a wavy, serrated edge.

37-year-old female presents with a headache


Jignesh Modi, MD




Axial T1 Pre-contrast



Axial T2 Pre-contrast



Axial T1
Post-contrast



Axial Gradient Echo


120



?


A large, stylized oak leaf graphic in a dark blue color, positioned on the left side of the slide, partially overlapping the title text.

Cavernous Malformation




Axial T1 Pre-contrast

Patchy increased T1 signal



Axial T2 Pre-contrast


Popcorn appearance



Axial T1
Post-contrast

Patchy increased
T1 signal

120



Axial Gradient
Echo

Blooming artifact

120

Cavernous Malfunction

- 40% are found incidentally on neuroimaging.
- Majority of symptomatic patients present at 40-60 years of age.
- Most often found as a single lesion.
- Multiple lesions may be familial; screening family members may be indicated when familial multiple cavernous malformation syndrome is suspected.

Imaging Features

CT: Unless large, these non-enhancing lesions are difficult to see on CT. If large, they appear as a region of hyperdensity resembling blood products and speckles of calcification. If there has been a recent hemorrhage, the lesion is more conspicuous and may be surrounded by a mantle of edema.

MRI is the modality of choice, demonstrating a characteristic “popcorn” or “berry” appearance with a rim of signal loss due to hemosiderin.

- **T1**

- Varied signal depending on the age of the blood products, small fluid-fluid levels may be evident

- **T2**

- Hypointense rim
- Varied signal internally depending on the age of blood products
- Blood locules with fluid-fluid levels may be seen
- If a recent bleed has occurred, surrounding edema may be present

GRE T2*/SWI

- Prominent blooming
- Useful for detecting smaller lesions otherwise missed by conventional spin echo sequences

T1 C+ (Gd): Usually no enhancement, although possible

Treatment & Prognosis

- Many cavernous malformations are asymptomatic and can be treated conservatively.
- Symptoms can relate to mass effect, epileptic activity or repeated hemorrhage. Symptomatic lesions should, when possible, be resected and complete resection is curative.

References

- Bergui M & Bradac G. Uncommon Symptomatic Cerebral Vascular Malformations. *AJNR Am J Neuroradiol.* 1997;18(4):779-83.
- Blitstein M & Tung G. MRI of Cerebral Microhemorrhages. *AJR Am J Roentgenol.* 2007;189(3):720-5. doi:10.2214/ajr.07.2249
- Jain R, Robertson P, Gandhi D, Gujar S, Muraszko K, Gebarski S. Radiation-Induced Cavernomas of the Brain. *AJNR Am J Neuroradiol.* 2005;26(5):1158-62. PMC8158583 - Pubmed
- Brunereau L, Labauge P, Tournier-Lasserre E, Laberge S, Levy C, Houtteville J. Familial Form of Intracranial Cavernous Angioma: MR Imaging Findings in 51 Families. *Radiology.* 2000;214(1):209-16. doi:10.1148/radiology.214.1.r00ja19209 - Pubmed
- Vilanova J, Barceló J, Smirniotopoulos J et al. Hemangioma from Head to Toe: MR Imaging with Pathologic Correlation. *RadioGraphics.* 2004;24(2):367-85. doi:10.1148/rg.242035079 - Pubmed
- Zhu W, Qi J, Zhan C et al. Magnetic Resonance Susceptibility Weighted Imaging in Detecting Intracranial Calcification and Hemorrhage. *Chin Med J (Engl).* 2008;121(20):2021-5. - Pubmed
- Pinker K, Stavrou I, Knosp E, Trattnig S. Are Cerebral Cavernomas Truly Nonenhancing Lesions and Thereby Distinguishable from Arteriovenous Malformations? *Magnetic Resonance Imaging.* 2006;24(5):631-7. doi:10.1016/j.mri.2005.10.037 - Pubmed
- Vogler R & Castillo M. Dural Cavernous Angioma: MR Features. *AJNR Am J Neuroradiol.* 1995;16(4):773-5. PMC8332321 - Pubmed
- Kondziolka D, Monaco III E, Lunsford L. Cavernous Malformations and Hemorrhage Risk. *Progress in Neurological Surgery.* 2012;27:141-6. doi:10.1159/000341774 - Pubmed
- Hegde A, Mohan S, Lim C. CNS Cavernous Haemangioma: “popcorn” in the Brain and Spinal Cord. *Clinical Radiology.* 2012;67(4):380-8. doi:10.1016/j.crad.2011.10.013 - Pubmed
- Dalyai R, Ghobrial G, Awad I et al. Management of Incidental Cavernous Malformations: A Review. *FOC.* 2011;31(6):E5. doi:10.3171/2011.9.focus11211
- Akers A, Al-Shahi Salman R, A Awad I et al. Synopsis of Guidelines for the Clinical Management of Cerebral Cavernous Malformations: Consensus Recommendations Based on Systematic Literature Review by