

56-year-old female with left-sided eye “bulge”

Joseph Ryan, MD, PhD

Gary Gong, MD, PhD

CT paranasal sinuses without IV contrast



CT paranasal sinuses without IV contrast



CT paranasal sinuses without IV contrast




CT paranasal sinuses without IV contrast




CT paranasal sinuses without IV contrast




CT paranasal sinuses without IV contrast



CT paranasal sinuses without IV contrast



CT paranasal sinuses without IV contrast



CT paranasal sinuses without IV contrast



A large, stylized oak leaf graphic in a dark blue color, positioned on the left side of the slide. It features detailed vein patterns and a lobed edge.

?

Orbital Lymphoma

Orbital Masses

- Broad spectrum of benign and malignant entities
- Categorized based on location and histology
- Cross-sectional imaging supplements clinical fundoscopic evaluation; imaging features reflect tissue composition
 - MRI gold standard; provides essential anatomic information about ocular structures involved, perineural spread, intracranial extension

Orbital Masses

- Vasculogenic lesions (17%), lymphoid-derived, lacrimal gland, optic nerve/meningeal, metastatic, peripheral nerve, primary melanoma (1%), and other rare ones: fibrocytic, myogenic, lipogenic, myxoid. (Compiled from Shields, et. al., case study)
- Anatomic compartment → possible tissue(s) of origin
- Evaluation of the bony orbit is crucial when orbital soft tissue mass is present
- Osseous remodeling (slow) vs. erosion (fast/aggressive)
- Bony foramina carry nerves → check for perineural spread, intracranial extension (optic canal, superior orbital fissure, infraorbital canal)

Orbit Anatomy

The orbit is divided into the intraconal space, extraconal space, muscle cone, ocular compartment

- Muscle cone → 5 of the 6 extraocular muscles (not the inferior oblique) converge at orbital apex to form tendinous ring (annulus of Zinn); divides the intra- and extra-conal spaces
- Ocular compartment/globe → continuous with CNS, comprises 3 layers (not typically visualized unless choroidal or retinal detachment)
 - Sclera – collagenous tissue layer continuous anteriorly with cornea, posteriorly with dura
 - Uvea – vascular pigmented layer, consists of iris, ciliary body, and choroid; *choroid very vascular, most frequent location for mets*
 - Retina – light sensory, continues posteriorly as optic nerve

Lymphoproliferative Lesions / Lymphoma

- Most common primary orbital tumor in adults >60 y/o. Subtypes include the following:
 - Lymphoid hyperplasia
 - Atypical lymphoid hyperplasia
 - Ocular adnexal (malignant) lymphoma
 - most common (67-90%), 24% of all space occupying orbital tumors in patients >60 years old
- Lesions may be 2/2 systemic lymphoma or primary
 - 30% with orbital lymphoma have systemic lymphoma within 10 years
 - MALT-subtype Non-Hodgkin most common

Orbital Lymphoma

- Typically present with painless proptosis, palpable mass, mildly restricted ocular motility.
 - Most (76%) unilateral, however if bilateral lesions in elderly, think lymphoma
 - 50% diffuse ill-defined, 50% smooth circumscribed
- T1-iso to muscle, T2-hyperintense to orbital fat; Uniform enhancement
- Tendency to mold to orbital structures, such as the globe, optic nerve, and orbital wall; tumor can cause bone remodeling, occasionally erosion
- Histology will show a range from benign reactive lymphoid hyperplasia, to highly mitotic atypical lymphocytes (typically B cell origin)

Orbital Lymphoma Mimics

Myositis Pseudotumor

- Idiopathic inflammation of rectus muscle
 - Usually presents with acute pain, periorbital edema
 - Responds rapidly to steroids
- Lymphoma usually pushes rectus, but not directly involved

Other mimics

- Granulomatosis with polyangiitis (Wegener's), IgG4-related disease (autoimmune), Erdheim-Chester disease (non-Langerhans histiocytosis)

References

- Knipe H. Orbital mass. Radiopaedia. 9/15/2022. <https://radiopaedia.org/articles/orbital-mass>
- Mosier, KM. Ill-defined Orbital Mass. STATdx. 7/10/2018.
- Davidson HC, Lantos JE. Orbital Lymphoproliferative Lesions. STATdx. 8/18/2021.
- Sarah N. Khan, MD; Ali R. Sepahdari, MD. Orbital masses: CT and MRI of common vascular lesions, benign tumors, and malignancies: Oculoplastic Imaging Update. Saudi Journal of Ophthalmology (2012) 26, 373–383.
- Tina D. Tailor MD, Divakar Gupta MD, Roberta W. Dalley MD, C. Dirk Keene MD PhD, Yoshimi Anzai MD MPH. Orbital Neoplasms in Adults: Clinical, Radiologic, and Pathologic Review. Head and Neck Neoplasms. RadioGraphics 2013; 33:1739–1758.