


31-year-old male with a history of seizures


Andrew Klufas, MD MBA
Racquel Helsing, MD

MR FLAIR




**UCONN
HEALTH
RADIOLOGY**

MR T2




UCONN
HEALTH
RADIOLOGY

MR T1




UCONN
HEALTH
RADIOLOGY

MR T1 C+



UCONN
HEALTH
RADIOLOGY

MR T2



**UCONN
HEALTH
RADIOLOGY**




?

UCONN
HEALTH
RADIOLOGY




Grey Matter Heterotopia

T2 FLAIR




T2



Periventricular, nodular ribbons of ectopic tissue
isointense to grey matter on all sequences


MR T2

Periventricular,
nodular ribbons of
ectopic tissue
isointense to grey
matter on all
sequences



T1

T1 MPR



Periventricular, nodular ribbons of ectopic tissue

isointense to grey matter on all sequences

UCONN
HEALTH
RADIOLOGY

Grey Matter Heterotopia

Interruption of the normal migration of neurons from the periventricular telencephalic germinal matrix to the cortex due to genetic abnormalities or infection/trauma

Divided macroscopically into

- Nodular heterotopias
 - Subependymal – most common
 - Subcortical
- Diffuse heterotopias
 - Band- also known as double cortex heterotopia and x-linked lissencephaly
 - Lissencephaly- types I and II
 - Laminar

Grey Matter Heterotopia

Clinical Presentation

- Most commonly present with focal seizures in the second decade of life
- +/- developmental delay or intellectual disability

Grey matter heterotopias are commonly associated with other congenital central nervous system anomalies including:

- Corpus callosum agenesis
- Schizencephaly
- Chiari II malformation
- Pachygryria, polymicrogyria
- Basilar cephaloceles

Imaging Findings

CT has limited use

- Difficult to appreciate small/focal involvement
- Slightly denser than surrounding white matter

MRI study of choice

- Heterotopic tissue follows same signal characters as grey matter on all sequences
- Also warrants careful examination to identify associated anomalies
- fMRI can demonstrate activation in heterotopic nodules, making epileptogenic EEG discharges

References

Gaillard F, Baba Y, Sharma R, et al. Grey matter heterotopia. Reference article, Radiopaedia.org (Accessed on 07 Sep 2023) <https://doi.org/10.53347/rID-1400>

Abdel Razek AA, Kandell AY, Elsorogy LG, Elmongy A, Basett AA. Disorders of cortical formation: MR imaging features. AJNR Am J Neuroradiol. 2009 Jan;30(1):4-11. doi: 10.3174/ajnr.A1223. Epub 2008 Aug 7. PMID: 18687750; PMCID: PMC7051699.

Barkovich AJ, Kuzniecky RI, Jackson GD, Guerrini R, Dobyns WB. A developmental and genetic classification for malformations of cortical development. Neurology. 2005 Dec 27;65(12):1873-87. doi: 10.1212/01.wnl.0000183747.05269.2d. Epub 2005 Sep 28. PMID: 16192428.

Leite CC, Lucato LT, Sato JR, Valente KD, Otaduy MC. Multivoxel proton MR spectroscopy in malformations of cortical development. AJNR Am J Neuroradiol. 2007 Jun-Jul;28(6):1071-5; discussion 1076-7. doi: 10.3174/ajnr.A0511. PMID: 17569960; PMCID: PMC8134160.

Kobayashi E, Bagshaw AP, Grova C, Gotman J, Dubeau F. Grey matter heterotopia: what EEG-fMRI can tell us about epileptogenicity of neuronal migration disorders. Brain. 2006 Feb;129(Pt 2):366-74. doi: 10.1093/brain/awh710. Epub 2005 Dec 9. PMID: 16339793.